IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

Stavenhagen, Jeffrey et al.

Appl. No. 10/754,922

Filed: January 9, 2004

For: Identification and Engineering of

Antibodies with Variant Fc Regions and Methods of Using

Same

Art Unit: **1643**

Examiner: Crowder, Chun

Atty. Docket: 1301.0004C

Confirmation No.: 8663

Supplemental Information Disclosure Statement Pursuant to 37 C.F.R. § 1.97(b)(1)

Honorable Commissioner for Patents P.O. Box 1450 Alexandria, VA 22313-1450

Sir:

Listed on accompanying substitute Form PTO SB08 are documents that may be considered material to the examination of this application, in compliance with the duty of disclosure requirements of 37 C.F.R. §§ 1.56, 1.97 and 1.98. Documents C01-C144 have been previously submitted, but are listed on the accompanying substitute Form PTO SB08 because their titles had not been previously provided to the Examiner. Copies of newly cited documents C145-C147 are submitted herewith.

- C145 Armstrong, S. *et al.* "Heterogeneity of IgG1 monoclonal anti-Rh(D): an investigation using ADCC and macrophage binding assays," Brit. J. Haematol. 66:257-262 (1987)
- C146 Kumpel, B.M. Brit. "Human monoclonal anti-D antibodies," J. Haematol. 71:415-420 (1989)
- Wiener, E. *et al.* "Differences between the activities of human monoclonal IgG1 and IgG3 anti-D antibodies of the Rh blood group system in their abilities to mediate effector functions of monocytes," Immunol. 65:159-163 (1988)

The Examiner is requested to contact the undersigned immediately in the event that any cited document may be unavailable, in order that a replacement copy can be provided.

The newly and previously submitted documents are:

Cite No.		cument Number atent Number	Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document
A01	US-	2003/0158389	08/21/03	Idusogie et al.
A02	US-	2004/0002587	1/1/04	Watkins et al.
A03	US-	2004/0110226	6/10/04	Lazar et al.
A04	US-	2004/0132101	7/8/04	Lazar et al.
A05	US-	2004/0185045	9/23/04	Koenig et al.
A06	US-	2005/0054832	3/10/05	Lazar et al.
A07	US-	4,752,601	06/21/88	Hahn
A08	US-	5,348,876	09/20/94	Michaelson et al.
A09	US-	5,576,184	11/19/96	Better et al.
A10	US-	5,585,089	12/17/96	Queen et al.
A11	US-	5,624,821	4/29/97	Winter et al.
A12	US-	5,648,260	07/15/97	Winter et al.
A13	US-	5,698,449	12/16/97	Baumann et al.
A14	US-	5,723,584	3/3/98	Schatz
A15	US-	5,736,135	4/7/98	Goeddel et al.
A16	US-	5,736,137	04/07/98	Anderson et al.
A17	US-	5,874,239	2/23/99	Schatz
A18	US-	5,932,433	8/3/99	Schatz
A19	US-	5,985,599	11/16/99	Mckenzie et al.
A20	US-	6,025,485	2/15/00	Kamb et al.
A21	US-	6,114,147	9/5/00	Frenken et al.
A22	US-	6,165,745	12/26/00	Ward et al.
A23	US-	6,194,551	2/27/01	Idusogie et al.
A24	US-	6,242,195	06/05/01	Idusogie et al.
A25	US-	6,277,375	08/21/01	Ward
A26	US-	6,300,065	10/9/01	Kieke et al.

In Re Appln of: Stavenhagen, Jeffrey US. Patent Appln. Serial No. 10/754,922

A27	US- 6,331,391	12/18/01	Wittrup et al.
A28	US- 6,423,538	7/23/02	Wittrup et al.
A29	US- 6,455,263	9/24/02	Payan
A30	US- 6,528,624	03/04/03	Idusogie et al.
A31	US- 6,538,124	03/25/03	Idusogie et al.
A32	US- 6,737,056	5/18/04	Presta
A33	US- 6,821,505	11/23/04	Ward
B01	EP 0 327 378	0809/89	Trustees of Columbia University
B02	WO 88/07089	09/22/88	Medical Research Council
B03	WO 89/07142	08/10/89	Morrison, S.
B04	WO 92/16562	10/01/92	Lynxvale Limited
B05	WO 93/22332	11/11/93	Board of Regents, Univ. Texas Sys
B06	WO 94/18330	8/18/94	Unilever PLC
B07	WO 94/29351	12/22/94	Morgan, S. et al.
B08	WO 95/05468	02/23/95	Lynxvale Limited
B09	WO 97/28267	08/07/97	Repligen Coroporation
B10	WO 97/34631	09/25/97	Board of Regents, Univ. Texas Sys
B11	WO 97/44362	11/27/97	Protein Design Labs, Inc.
B12	WO 98/05787	02/12/98	Bristol-Myers Squibb Company
B13	WO 98/23289	06/04/98	The General Hospital Corporation
B14	WO 98/52975	11/26/98	Duetches Krebsforschungzentrum Stiftung Des Offentlichen Rechts
B15	WO 99/43713	09/02/99	Lexigen Pharmaceuticals Corp.
B16	WO 99/51642	10/14/99	Genentech, Inc
B17	WO 99/58572	11/18/99	Cambridge Univ. Tech. Serv. Ltd.
B18	WO 00/09560	02/24/00	Abgenix, Inc.
B19	WO 00/42072	7/20/00	Genenech, Inc.
B20	WO 02/060919	08/0802	Medimmune, Inc.
B21	WO 02/086070	10/31/02	Dyax Corporation, Inc.
B22	WO 03/074679	09/12/03	Xencor
B23	WO 04/029207	04/08/04	Xencor
B24	WO 04/074455	09/02/04	Applied Molecular Evolution, Inc.
B25	WO 04/099249	11/18/04	Xencor

B26	WO 05/070963	08/04/05	Applied Molecular Evolution, Inc.
B27	WO 06/020114	02/23/06	Applied Molecular Evolution, Inc.
C01	Altman <i>et al.</i> , Pheno 274:94-96, 1996	otypic Analysis of An	tigen-Specific T Lymphocytes", Science
C02			ution abolishes the heterogeneity of chimeric munol 30:105-108, 1993
C03			ing interactions of human and herpes simplex Transactions 30:495-500, 2002
C04			molecules lacking Fegamma receptor I s," Eur J Immunol 29:2613-2624, 1999
C05			man FcgammaRIIa and FcgammaRIIb tant antibodies," Mol Immunol 40:585-593,
C145			G1 monoclonal anti-Rh(D): an investigation says," Brit. J. Haematol. 66:257-262 (1987)
C06			ls for the detection and evaluation of drugs Experientia. Oct 15;44(10):841-848, 1988
C07		1997, "Yeast surface s", Nature Biotechnol	display for screening combinatorial ogy 15:553-557
C08	Boder and Wittrup, Giotechnol Prog 14:		f surface-displayed polypeptide libraries,"
C09			y for directed evolution of protein expression, nology 328:430-444, 2000
C10			ody fragments with monovalent femtomolar d. Sci. USA 97:10701-10705, 2000
C11	polymorphic forms i		nma RIIa (CD32) and Fc gamma RIIIb (CD16) man IgG1- and IgG3-opsonized bacteria and 1994
C12			fic amino acid residues affecting complementur J Immunol 24:2542-2547, 1994
C13		(Microbes as Tools fo	or Cell Biology) in <i>Methods In Cell Biololgy</i> , 64, 1994
C14	Burlmeister et al., "Ger" Fc," Nature 372:379		e complex of rat neonatal Fc receptor with
C15	Burton and Woof, "I 84, 1992	Human antibody effe	ctor function," Advances in Immunology 51:1-
C16	Burton et al., "Mole Mol Immunol 25:11	_	entibody (IgG) by cellular Fc receptor (FcRI),"
C17	Burton, "Immunogle	obulin G: functional s	ites," Mol Immunol 22:161-206, 1985

C18	Canfield and Morrison, "The binding affinity of human IgG for its high affinity Fc receptor is determined by multiple amino acids in the CH2 domain and is modulated by the hinge region," J Exp Med 173:1483-1491, 1991
C19	Caron et al., "Engineered humanized dimeric forms of IgG are more effective antibodies," J Exp Med 176:1191-5, 1992
C20	Carter <i>et al.</i> , "Humanization of an anti-p185HER2 antibody for human cancer therapy," Proc. Natl. Acad. Sci. USA 89:4285-4289, 1992
C21	Cartron et al., "Therapeutic activity of humanized anti-CD20 monoclonal antibody and polymorphism in IgG Fc receptor FcgammaRIIIa gene," Blood 99:754-758, 2002
C22	Chappel <i>et al.</i> , "Identification of the Fc gamma receptor class I binding site in human IgG through the use of recombinant IgG1/IgG2 hybrid and point-mutated antibodies," Proc. Natl. Acad. Sci USA 88:9036-9040, 1991
C23	Chappel et al., "Identification of a secondary Fc gamma RI binding site within a genetically engineered human IgG antibody," J Biol. Chem 268:25124-25131, 1993
C24	Ciccimarra <i>et al.</i> , "Localization of the IgG effector site for monocyte receptors," Proc. Natl. Acad. Sci. U.S.A. 72:2081-2083, 1975
C25	Clynes and Ravetch, "Cytotoxic antibodies trigger inflammation through Fc receptors," Immunity 3:21-26, 1995
C26	Clynes <i>et al.</i> , "Modulation of immune complex-induced inflammation in vivo by the coordinate expression of activation and inhibitory Fc receptors," J Exp Med 189:179-185, 1999
C27	Clynes et al., "Inhibitory Fc receptors modulate in vivo cytoxicity against tumor targets," Nature Medicine 6:443-446, 2000
C28	Clynes <i>et al.</i> , "Fc receptors are required in passive and active immunity to melanoma," Proc. Natl. Acad. Sci USA 95:652-656, 1998
C29	Clynes et al., "Uncoupling of immune complex formation and kidney damage in autoimmune glomerulonephritis," Science 279:1052-1054, 1998
C30	de Haas, Wien Kin "IgG-Fc receptors and the clinical relevance of their polymorphisms," Wien Klin Wochenscha 113:825-831, 2001
C31	Deisenhofer, "Crystallographic refinement and atomic models of a human Fc fragment and its complex with fragment B of protein A from Staphylococcus aureus at 2.9- and 2.8-A resolution," Biochem. 20:2361-2370, 1981
C32	Deo et al., "Clinical significance of IgG Fc receptors and Fc gamma R-directed immunotherapies," Immunology Today 18:127-135, 1997
C33	Duncan and Winter, "The binding site for C1q on IgG," Nature 332:738-740, 1988
C34	Duncan and Winter, "Localization of the binding site for the human high-affinity Fc receptor on IgG," Nature 332:563-564, 1988
C35	Flesch and Neppert, "Functions of the Fc receptors for immunoglobulin G," J Clin Lab Anal 14:141-156, 2000
C36	Gergeley et al., "Fc receptors on lymphocytes and K cells," Biochemical Society Transactions 12:739-743, 1984

C37	Gergely and Sarmay, "The two binding-site models of human IgG binding Fc gamma receptors," FASEB J 4:3275-3283, 1990
C38	Greenwood and Clark, Effector functions of matched sets of recombinant human IgG subclass antibodies". (final version edited Feb. 11, 1993)
C39	Greenwood <i>et al.</i> , "Structural motifs involved in human IgG antibody effector functions," Eur J Immunol 23:1098-1104, 1993
C40	Greenwood <i>et al.</i> , "Engineering multiple-domain forms of the therapeutic antibody CAMPATH-1H: effects on complement lysis," Therapeutic Immunology 1:247-255, 1994
C41	Hadley <i>et al.</i> , "The functional activity of Fc gamma RII and Fc gamma RIII on subsets of human lymphocytes," Immunology 76:446-451, 1992
C42	Hatta et al., "Association of Fc gamma receptor IIIB, but not of Fc gamma receptor IIA and IIIA polymorphisms with systemic lupus erythematosus in Japanese," Genes and Immunity 1:53-60, 1999
C43	Hayes, Fc Engineering to Enhance Monoclonal Antibody Effector Functions. (Presentation) Xecor, CA, 2003
C44	Herzenberg <i>et al.</i> , "The history and future of the fluorescence activated cell sorter and flow cytometry: a view from Stanford," Clinical Chem. 2002;48:1819-1827, 2002
C45	Heyman, "Regulation of antibody responses via antibodies, complement, and Fc receptors," Annu Rev Immunol 18:709-737, 2000
C46	Hogarth et al., "Characterization of FcR Ig-binding sites and epitope mapping," Immunomethods 4:17-24, 1994
C47	Holler <i>et al.</i> , "In vitro evolution of a T cell receptor with high affinity for peptide/MHC," Proc. Natl. Acad. Sci. U.S.A. 97:5387-92, 2000
C48	Hulett <i>et al.</i> , "Identification of the IgG binding site of the human low affinity receptor for IgG Fc gamma RII. Enhancement and ablation of binding by site-directed mutagenesis," J. Biol. Chem. 269:15287-15293, 1994
C49	Hulett et al., "Multiple regions of human Fc gamma RII (CD32) contribute to the binding of IgG," J. Biol. Chem. 270:21188-21194, 1995
C50	Hulett et al., "Chimeric Fc receptors identify functional domains of the murine high affinity receptor for IgG," J Immunol 147:1863-1868, 1991
C51	Idusogie et al., "Mapping of the C1q binding site on rituxan, a chimeric antibody with a human IgG1 Fc," J Immunol 164: 4178-4184, 2000
C52	Idusogie <i>et al.</i> , "Engineered antibodies with increased activity to recruit complement," J Immunol 166:2571-2575, 2001
C53	Isaacs <i>et al.</i> , "A therapeutic human IgG4 monoclonal antibody that depletes target cells in humans," Clin Exp Immunol 106 :427-433, 1996
C54	Isaacs et al., "Therapy with monoclonal antibodies. An in vivo model for the assessment of therapeutic potential," J Immunol 148:3062-3071, 1992

- Isaacs *et al.*, "Therapy with monoclonal antibodies. II. The contribution of Fc gamma receptor binding and the influence of C(H)1 and C(H)3 domains on in vivo effector function," J Immunol 161:3862-3869, 1998
- C56 Jassal *et al.*, "Remodeling glycans on IgG by genetic re-engineering," Biochem Soc Trans 26:S113, 1998
- C57 Jefferis and Lund, "Interaction sites on human IgG-Fc for FcgammaR: current models," Immunology Letters 82:57-65, 2002
- C58 Jefferis *et al.*, "Recognition sites on human IgG for Fc gamma receptors: the role of glycosylation," Immunol Lett 44:111-7, 1995
- C59 Jefferis *et al.*, "IgG-Fc-mediated effector functions: molecular definition of interaction sites for effector ligands and the role of glycosylation," Immunol Rev 163:59-76, 1998
- C60 Jefferis *et al.*, "Molecular definition of interaction sites on human IgG for Fc receptors (huFc gamma R)," Mol Immunol 27:1237-1240, 1990
- C61 Jendeberg *et al.*, "Engineering of Fc(1) and Fc(3) from human immunoglobulin G to analyse subclass specificity for staphylococcal protein A," J Immunological Methods 201:25-34, 1997
- C62 Kadar *et al.*, "Synthetic peptides comprising defined sequences of CH-2 and CH-3 domains of human IgG1 induce prostaglandin E2 production from human peripheral blood mononuclear cells," Immunol Lett 32:59-63, 1992
- C63 Kadar *et al.*, "Modulatory effect of synthetic human IgG Fc peptides on the in vitro immune response of murine spleen cells," Int J Immunpharmacol 13:1147-55, 1991
- C64 Kato et al., "Structural basis of the interaction between IgG and Fcγ receptors," J Mol Biol 295:213-224, 2000
- Keler *et al.*, "Differential effect of cytokine treatment on Fc alpha receptor I- and Fc gamma receptor I-mediated tumor cytotoxicity by monocyte-derived macrophages," J. of Immunol. 164:5746-52, 2000
- Kieke *et al.*, "Selection of functional T cell receptor mutants from a yeast surface-display library," Proc. Natl. Acad. Sci. U.S.A. 96:5651-56, 1999
- Kim *et al.*, "Analysis of FcγRIII and IgG Fc polymorphism reveals functional and evolutionary implications of protein-protein interaction," J Mol Evol 53:1-9, 2001
- C68 Klein *et al.*, "Expression of biological effector functions by immunoglobulin G molecules lacking the hinge region," Proc. Natl. Acad. Sci. U.S.A. 78:524-528, 1981
- Koene *et al.*, "Fc gammaRIIIa-158V/F polymorphism influences the binding of IgG by natural killer cell Fc gammaRIIIa, independently of the Fc gammaRIIIa-48L/R/H phenotype," Blood 90:1109-1114, 1997
- Kranz *et al.*, "Mechanisms of ligand binding by monoclonal anti-fluorescyl antibodies,"J. Biol. Chem. 257:6987-6995, 1982
- C146 Kumpel, B.M. Brit. "Human monoclonal anti-D antibodies," J. Haematol. 71:415-420 (1989)
- C71 Lehmann *et al.*, "Phagocytosis: measurement by flow cytometry," J Immunol Methods. 243(1-2):229-42, 2000

C72	Lehrnbecher et al., "Variant genotypes of the low-affinity Fcgamma receptors in two
	control populations and a review of low-affinity Fegamma receptor polymorphisms in
	control and disease populations," Blood 94:4220-4232, 1999

- C73 Li *et al.*, "Reconstitution of human Fc gamma RIII cell type specificity in transgenic mice," J Exp Med 183:1259-1263, 1996
- C74 Liu *et al.*, "Production of a mouse-human chimeric monoclonal antibody to CD20 with potent Fc-dependent biologic activity," J. Immunol. 139:3521-3526, 1987
- C75 Lund *et al.*, "Expression and characterization of truncated forms of humanized L243 IgG1. Architectural features can influence synthesis of its oligosaccharide chains and affect superoxide production triggered through human Fegamma receptor I," Eur J Biochem 267:7246-57, 2000
- C76 Lund *et al.*, "Oligosaccharide-protein interactions in IgG can modulate recognition by Fc gamma receptors," FASEB J 9:115-119, 1995
- C77 Lund *et al.*, "Human Fc gamma RI and Fc gamma RII interact with distinct but overlapping sites on human IgG," J Immunol 147:2657-62, 1991
- C78 Lund *et al.*, "Multiple interactions of IgG with its core oligosaccharide can modulate recognition by complement and human Fc gamma receptor I and influence the synthesis of its oligosaccharide chains," J Immunol 157:4963-4969, 1996
- C79 Lund *et al.*, "Multiple binding sites on the CH2 domain of IgG for mouse Fc gamma R11," Molecular Immunology 29:53-59, 1992
- C80 Maenaka *et al.*, "The human low affinity Fegamma receptors IIa, IIb, and III bind IgG with fast kinetics and distinct thermodynamic properties," J Biol Chem 48:44898-904, 2001
- C81 Michaelsen *et al.*, "One disulfide bond in front of the second heavy chain constant region is necessary and sufficient for effector functions of human IgG3 without a genetic hinge," Immunolgy 91:9243-9247, 1994
- C82 Morgan *et al.*, "The N-terminal end of the CH2 domain of chimeric human IgG1 anti-HLA-DR is necessary for C1q, Fc gamma RI and Fc gamma RIII binding," Immunology 86:319-324, 1995
- C83 Morrison *et al.*, "Structural determinants of IgG structure," Immunologist 2:119-124, 1994
- Munn *et al.*, "Phagocytosis of tumor cells by human monocytes cultured in recombinant macrophage colony-stimulating factor," J Exp Med. 172(1):231-7, 1990
- Nagarajan *et al.*, "Ligand binding and phagocytosis by CD16 (Fc gamma receptor III) isoforms. Phagocytic signaling by associated zeta and gamma subunits in Chinese hamster ovary cells," J Biol Chem 270:25762-25770, 1995
- Neuberger *et al.*, "Recombinant antibodies possessing novel effector functions," Nature 312:604-608, 1984
- Norderhaug *et al.*, "Chimeric mouse human IgG3 antibodies with an IgG4-like hinge region induce complement-mediated lysis more efficiently than IgG3 with normal hinge," Eur J Immunol 21:2379-84, 1991
- C88 Nose and Leanderson, "Substitution of asparagine 324 with aspartic acid in the Fc

	portion of mouse antibodies reduces their capacity for C1q binding," Eur J Immunol 19:2179-81, 1989
C89	Okazaki <i>et al.</i> , "Fucose depletion from human IgG1 oligosaccharide enhances binding enthalpy and association rate between IgG1 and FcgammaRIIIa," J Mol Biol 336:1239-1249, 2004
C90	Orfao and Ruiz-Arguelles, "General concepts about cell sorting techniques," Clinical Biochem. 29:5-9, 1996
C91	Partridge et al., "The use of anti-IgG monoclonal antibodies in mapping the monocyte receptor site on IgG," Mol Immunol. 23(12):1365-72, 1986
C92	Perussia "Human Natural Killer Cell Protocols" in <i>Methods Molecular Biology</i> . vol. 121 (Campbell <i>et al</i> . eds.) Humana Press Inc., Totowa, NJ. 179-92, 2000
C93	Radaev and Sun, "Recognition of immunoglobulins by Fcgamma receptors," Molecular Immunology 38:1073-1083, 2001
C94	Ravetch and Bolland, "IgG Fc receptors," Annu Rev Immunol 19:275-90, 2001
C95	Ravetch and Clynes, "Divergent roles for Fc receptors and complement in vivo," Annu Rev Immunol 16:421-432, 1998
C96	Ravetch and Kinet, "Fc receptors," Annu Rev Immunol 9:457-492, 1991
C97	Ravetech and Lanier, "Immune inhibitory receptors," Science 290:84-89, 2000
C98	Redpath <i>et al.</i> , "The influence of the hinge region length in binding of human IgG to human Fegamma receptors," Hum Immunol 59:720-727, 1998
C99	Reff et al., "Depletion of B cells in vivo by a chimeric mouse human monoclonal antibody to CD20," Blood 83:435-445, 1994
C100	Riechmann et al., "Reshaping human antibodies for therapy," Nature. 332(6162):323-7, 1988
C101	Sarmay <i>et al.</i> , "The effect of synthetic peptides corresponding to Fc sequences in human IgG1 on various steps in the B cell activation pathway," Eur J Immunol 18:289-294, 1988
C102	Sarmay et al., "Ligand inhibition studies on the role of Fc receptors in antibody-dependent cell-mediated cytotoxicity," Mol Immunol 21:43-51, 1984
C103	Sarmay et al., "Mapping and comparison of the interaction sites on the Fc region of IgG responsible for triggering antibody dependent cellular cytotoxicity (ADCC) through different types of human Fc gamma receptor," Mol Immunol 29:633-639, 1992
C104	Sautes-Fridman <i>et al.</i> , "Fc gamma receptors: a magic link with the outside world," ASHI Quarterley, 4 th Quarter:148-151, 2003
C105	Schaffner <i>et al.</i> , "Chimeric interleukin 2 receptor alpha chain antibody derivatives with fused mu and gamma chains permit improved recruitment of effector functions," Mol Immunol 32:9-20, 1995 (Erratum in 32:1299, 1995)
C106	Schatz <i>et al.</i> , "Use of peptide libraries to map the substrate specificity of a peptide-modifying enzyme: a 13 residue consensus peptide specifies biotinylation in Escherichia coli," Bio/Technology 11:1138-1143, 2000

C123

C107	Sensel <i>et al.</i> , "Amino acid differences in the N-terminus of C(H)2 influence the relative abilities of IgG2 and IgG3 to activate complement," Molecular Immunology 34:1019-1029, 1997
C108	Shields <i>et al.</i> , "High resolution mapping of the binding site on human IgG1 for Fc gamma RI, Fc gamma RII, Fc gamma RIII, and FcRn and design of IgG1 variants with improved binding to the Fc gamma R," J Biol Chem 276:6591-6604, 2001
C109	Shopes <i>et al.</i> , "Recombinant human IgG1-murine IgE chimeric Ig. Construction, expression, and binding to human Fc gamma receptors," J Immunol 145:3842-3848, 1990
C110	Shopes, "A genetically engineered human IgG mutant with enhanced cytolytic activity," J Immunol 148:2918-2922, 1992
C111	Shopes, "A genetically engineered human IgG with limited flexibility fully initiates cytolysis via complement," Molecular Immunology 30:603-609, 1993
C112	Shusta et al., "Yeast polypeptide fusion surface display levels predict thermal stability and soluble secretion efficiency," J Mol Biol 292:949-956, 1999
C113	Shusta <i>et al.</i> , "Increasing the secretory capacity of Saccharomyces cerevisiae for production of single-chain antibody fragments," Nature Biotechnology 16:773-777, 1998
C114	Shusta <i>et al.</i> , "Directed evolution of a stable scaffold for T-cell receptor engineering," Nature Biotechnology 18:754-759, 2000
C115	Smith and Morrison, "Recombinant polymeric IgG: an approach to engineering more potent antibodies," Bio/Technology 12:683-688, 1994
C116	Sondermann and Oosthuizen, "The structure of Fc receptor/Ig complexes: considerations on stoichiometry and potential inhibitors," Immunology Letters, 82:51-56, 2002
C117	Sondermann <i>et al.</i> , "Molecular basis for immune complex recognition: a comparison of Fc-receptor structures," J. Mol. Biol. 309:737-749, 2001
C118	Sondermann <i>et al.</i> , "Crystal structure of the soluble form of the human fcgamma-receptor IIb: a new member of the immunoglobulin superfamily at 1.7 A resolution," EMBO J 18:1095-1103, 1999
C119	Sondermann <i>et al.</i> , "The 3.2-A crystal structure of the human IgG1 Fc fragment-Fc gammaRIII complex," Nature 406:267-273, 2000
C120	Steplewski <i>et al.</i> , "Biological activity of human-mouse IgG1, IgG2, IgG3, and IgG4 chimeric monoclonal antibodies with antitumor specificity," Proc. Natl. Acad. Sci. U.S.A. 85:4852-4856, 1988
C121	Strohmeier <i>et al.</i> , "Role of the Fc gamma R subclasses Fc gamma RII and Fc gamma RIII in the activation of human neutrophils by low and high valency immune complexes," J Leukocyte Biol 58:415-422, 1995
C122	Sylvestre and Ravetch, "A dominant role for mast cell Fc receptors in the Arthus reaction," Immunity 5:387-390, 1996

Sylvestre and Ravetch, "Fc receptors initiate the Arthus reaction: redefining the

inflammatory cascade," Science 265:1095-1098, 1994

C124	Takai <i>et al.</i> , "FcR gamma chain deletion results in pleiotrophic effector cell defects," Cell 76:519-529, 1994
C125	Takai et al., "Augmented humoral and anaphylactic responses in Fc gamma RII-deficient mice," Nature 379:346-349, 1996
C126	Takai, "Roles of Fc receptors in autoimmunity," Nature Reviews 2:580-592, 2002
C127	Tamm et al., "The IgG binding site of human FcγRIIIB receptor involves CC' and FG loops of the membrane-proximal domain," J Biol Chem 271:3659-3666, 1996
C128	Tao et al., "The differential ability of human IgG1 and IgG4 to activate complement is determined by the COOH-terminal sequence of the CH2 domain," J Exp Med 173:1025-1028, 1991
C129	Tao et al., "Structural features of human immunoglobulin G that determine isotype-specific differences in complement activation," J Exp Med 178:661-667, 1993
C130	Van Sorge <i>et al.</i> , "FcgammaR polymorphisms: Implications for function, disease susceptibility and immunotherapy," Tissue Antigens 61:189-202, 2003
C131	VanAntwerp and Wittrup, "Fine affinity discrimination by yeast surface display and flow cytometry," Biotechnol Prog 16:31-37, 2000
C132	Vidarte, "Serine 132 is the C3 covalent attachment point on the CH1 domain of human IgG1," J Biol Chem 276:38217-38233, 2001
C133	Ward and Ghetie, "The effector functions of immunoglobulins: implications for therapy," Therapeutic Immunology 2:77-94, 1995
C134	Weng and Levy, "Two immunoglobulin G fragment C receptor polymorphisms independently predict response to rituximab in patients with follicular lymphoma," J Clin Oncol 21:3940-3947, 2003
C147	Wiener, E. <i>et al</i> . "Differences between the activities of human monoclonal IgG1 and IgG3 anti-D antibodies of the Rh blood group system in their abilities to mediate effector functions of monocytes," Immunol. 65:159-163 (1988)
C135	Wing <i>et al.</i> , "Mechanism of first-dose cytokine-release syndrome by CAMPATH 1-H: Involvement of CD16 (FcγRIII) and CD11a/CD18 (LFA-1) on NK cells," J Clin Invest 98:2819-2826, 1996
C136	Wingren <i>et al.</i> , "Comparison of surface properties of human IgA, IgE, IgG and IgM antibodies with identical and different specificities," Scand J Immunol 44:430-436, 1996
C137	Wittrup, "The single cell as a microplate well," Nat Biotechnol 18:1039-1040, 2000
C138	Witttrup, "Protein engineering by cell-surface display," Curr, Opin. Biotechnol. 12:395-399, 2001
C139	Woof <i>et al.</i> , "Localisation of the monocyte-binding region on human immunoglobulin G," Mol Immunol 23:319-330, 1986
C140	Wu <i>et al.</i> , "a novel polymorphism of FcγRIIIa (CD16) alters receptor function and predisposes to autoimmune disease," J Clin Invst 100:1059-1070, 1997
C141	Xu et al., "Residue at position 331 in the IgG1 and IgG4 CH2 domains contributes to

their differential ability to bind and activate complement," J Biol Chem 269:3469-

58 :3905-3908, 1998

Yeung and Wittrup, "Quantitative screening of yeast surface-displayed polypeptide libraries by magnetic bead capture," Biotechnol Prog 18:212-220, 2002
Zeidler et al., "The Fc-region of a new class of intact bispecific antibody mediates activation of accessory cells and NK cells and induces direct phagocytosis of tumour cells," British J Cancer 83:261-266, 2000
Zuckier et al., "Chimeric human-mouse IgG antibodies with shuffled constant region exons demonstrate that multiple domains contribute to in vivo half-life," Cancer Res

This Information Disclosure Statements is being submitted prior to an Initial Office Action. No fee is accordingly believed due for consideration of this Information Disclosure Statement. However, if the Commissioner determines that an additional fee is required in for consideration of this Information Disclosure Statement, the U.S. Patent and Trademark Office is hereby authorized to charge any fee deficiency, or credit any overpayment, to Deposit Account No. **05-0460** referencing docket number **0301.0004C**.

The submission of the listed and appended documents is not intended as an admission that any such document constitutes prior art against the claims of the present application. Applicant does not waive any right to take any action that would be appropriate to antedate or otherwise remove any listed document as a competent reference against the claims of the present application.

Applicant respectfully requests that the documents listed on the accompanying Form SB08 be considered and made of record in the present application. Applicant further requests that the Examiner initial and return a copy of the enclosed PTO SB08 form and indicate in the official file wrapper of this patent application that the documents have been considered.

- 13 -

While the listed references are considered relevant to the prosecution of the present application, it is submitted that the references, either alone or in combination, do not detract from the patentability of the claimed invention.

Respectfully Submitted,

Date: <u>December 7, 2006</u> Edell, Shapiro & Finnan, LLC 1901 Research Blvd., Suite 400

Rockville, MD 20850

Telephone: (301) 424-3640 Facsimile: (301) 762-4056 /Jeffrey I. Auerbach/

Jeffrey I. Auerbach Registration No. 32,680 Attorney for Assignee